


# Fault geometry used in RSQ









# Strike-slip events

Raw data from RSQsim:

1. M
2. SR
3. A
4. L (FSL, FML ...)
5. AD

Derived data from them:

1. W (A/L)
2. Aspect ratio (L/W)
3. SDP (stress drop from AD, L and W)



# Comparison of subgroups (fault maturity)

Immature fault:


Intermediate mature fault:

Mature fault:

$0.01 < SR < 1$  mm/yr



$1 < SR < 10$  mm/yr

$10 < SR < 40$  mm/yr



# Stress drop vs geological slip rate


$$\begin{aligned} \text{logsigma} &= 1.68 + -0.22 * \text{log}(\text{SR}) \\ \text{logsigma} &= 0.71 + -0.23 * \text{log}(\text{SR}) \\ \text{logsigma} &= 0.78 + -0.10 * \text{log}(\text{SR}) \\ \text{logsigma} &= 0.85 + -0.08 * \text{log}(\text{SR}) \end{aligned}$$




# Reverse and normal events

Reverse


normal




Reverse









normal













# Fault style dependence







# Change definition of L



# Next validation?

## Geophysical Research Letters

RESEARCH LETTER  
10.1029/2021GL095271



### Fault Interactions Enhance High-Frequency Earthquake Radiation

Shanna X. Chu<sup>1</sup>, Victor C. Tsai<sup>1</sup>, Daniel T. Trugman<sup>2</sup>, and Greg Hirth<sup>1</sup>

<sup>1</sup>Department of Earth, Environmental and Planetary Sciences, Brown University, Providence, RI, USA, <sup>2</sup>Jackson School of Geosciences, The University of Texas at Austin, Austin, TX, USA





#### Key Points:

- We introduce two new metrics quantifying the complexity of interacting faults
- In Southern California, enhanced



A recent study connects stress drop of seismicity with the fault surface trace complexity and a power law has been found.

(a) misalignment,  $R_M$





# Evolving geometrical heterogeneities of fault trace data

Neta Wechsler, Yehuda Ben-Zion and Shari Christofferson

Department of Earth Sciences, University of Southern California, Los Angeles, CA 90089-0740, USA. E-mail: wechsler@usc.edu

Accepted 2010 April 28. Received 2010 March 15; in original form 2009 November 11



ifornia with the faults used in this study. The fault zones are marked by their defining rectangles. See

**Figure 7.** (a) The relation between the circular standard deviation, the misalignment from plate motion direction and the recent slip rate for each fault zone. Black line is the least squares fit for the data using a log-linear relation. The grey line is the same, but dropping the Rinconada data point. The corresponding equations and  $R^2$  values are next to each fit. (b) Same as a, but for the circular standard error.

Can we derive a relation of SR-sigma (surface trace complexity) from seismicity (may or may not support that from previous study)? Any thoughts?

